

INDUSTRIAL PHARMACY

1. The following tablet is intended to be added to given volume of water by pharmacist to obtain certain drug concentration before administration.

- (A) Hypodermic tablet
- (B) Dispensing tablet
- (C) Efferevescent tablet
- (D) Tablet triturates

2. Chilsonator is used in which technique of tablet preparation.

- (A) Direct compression
- (B) Dry granulation
- (C) Wet granulation
- (D) None

3. The mechanical strength of the tablet can be found by

- (A) Friability
- (B) Disintegration
- (C) Dissolution
- (D) Hardness

4. Starch paste is used in tablet manufacturing as

- (A) Granulating agent
- (B) Disintegrating agent
- (C) Diluent
- (D) Lubricant

5. Which of the following excipient is hydrolysed starch.

- (A) Sugartab
- (B) Nutab
- (C) Celutab
- (D) All

6. The relative humidity and temperature to be maintained during dry granulation of Aspirin tablet.

- (A) <30% and 70°F
- (B) >30% and 70°F
- (C) >30%and 70°C
- (D) >40%and 60°F

7. Which of the following is both wet and dry granulating mixer (high speed mixer)

- (A) Sigma
- (B) Planetary
- (C) Both A and B
- (D) Diosna mixer



YOUR SUCCESS IS OUR MOTTO

8. Hardness can be tested by the testers

- (A) Pfizer
- (B) Erweka
- (C) Schleuniger
- (D) All the above

9. Cerelose is brand name of

- (A) Sucrose
- (B) Dextrose
- (C) Cellulose
- (D) Starch

10. Ac-Di-Sol is a

- (A) cross linked MCC
- (B) cross linked PVP
- (C) cross linked sodium CMC
- (D) cross linked sodium starch glycolate

11. The disintegration time of sugar coated tablet is

- (A) NMT 15 min
- (B) NMT 30 min
- (C) NMT 45min
- (D) NMT 60min

12. Which of the following diluent has negative heat of solution

- (A) Sucrose
- (B) Manitol
- (C) Sorbitol
- (D) Glucose

13. According to I.P., % difference permitted in weight variation test of 240 mg tablet.


- (A) $\pm 1\%$
- (B) $\pm 5\%$
- (C) $\pm 7.5\%$
- (D) $\pm 10\%$

14. Air entrapment in tablet among granules can be overcome by

- (A) Pre-compression
- (B) Reducing final compression force
- (C) Minimising tableting rate
- (D) All the above

15. The temp maintained in dissolution apparatus is

- (A) $37 \pm 0.5^\circ \text{C}$



YOUR SUCCESS IS OUR MOTTO

MYBO's GPAT/ NIPER/ PGCET BOOSTER PACK-3

(From MYBO Knowledge Center) : MYBO – A Voluntary Organization

- (B) $37 \pm 1^\circ \text{C}$
- (C) $37 \pm 1.5^\circ \text{C}$
- (D) $37 \pm 2^\circ \text{C}$

MYBO's GPAT/ NIPER/ PGCET BOOSTER PACK-3 is provided by

MYBO Team member:

Ms. AAYISHA BEGUM, (M.Pharm),

Andhra University College of Pharmaceutical Sciences, Visakhapatnam.

NOTE:

1. Key for the above bits is available with MYBO's GPAT/ NIPER/ PGCET BOOSTER PACK-4
2. If you are interested to help the GPAT/ NIPER/ PGCET Aspirants, you can send 15 bits each with four options (correct answer underlined) to mybogroup@gmail.com After verification, we will publish them on your name at www.mybogroup.com



YOUR SUCCESS IS OUR MOTTO

KEY @ MYBO's BOOSTER PACK-2

- | | | | | |
|-------|-------|-------|-------|-------|
| 1. B | 2. C | 3. A | 4. C | 5. C |
| 6. A | 7. D | 8. C | 9. B | 10. D |
| 11. A | 12. B | 13. A | 14. B | 15. B |