


PHARMACOLOGY


- 1. Which of the below drug causes diabetes**
 - (A) Diazoxide
 - (B) Triamterene
 - (C) Digoxin
 - (D) Phenacetin
- 2. Interaction between chlorpromazine and levo carbidopa causes**
 - (A) Anti-parkinson effect
 - (B) Antagonistic effect
 - (C) Lymphoma
 - (D) None
- 3. Pin point pupil is a characteristic feature of**
 - (A) Morphine abusers
 - (B) Phenytoin
 - (C) Phenacetin
 - (D) Minoxidil
- 4. Accumulation of beta amyloid plaques in various areas of brain like cortex is characteristically seen in**
 - (A) Alzheimers disease
 - (B) Non-hogdkins lymphoma
 - (C) Meningitis
 - (D) All the above
- 5. Ketoconazole + coca cola drink, results in the interaction of**
 - (A) Increased systemic absorption
 - (B) Decreased systemic absorption
 - (C) No effect on absorption
 - (D) None
- 6. Wilsons disease is resulted due to a metabolic defect of the element**
 - (A) Copper
 - (B) Iron
 - (C) Cyanide
 - (D) Gold
- 7. Which of the following is highly teratogenic drug**
 - (A) Omeprazole
 - (B) Ranitidine



MYBO's GPAT/ NIPER/ PGE CET BOOSTER PACK-8

(From MYBO Knowledge Center) : MYBO – A Voluntary Organization

- (C) Isotretinoin
(D) Nifedipine
8. **The currently used most efficacious drug for scabies in pediculosis which shows 100% cure rate**
(A) Lindane
(B) Permethrin
(C) Naftitine
(D) Crotamiton
9. **Selective oestrogen receptor down regulator is**
(A) Letrazole
(B) Toremifone
(C) Dutasteride
(D) Fulvestrant
10. **Neural tube defects occurs with one of the anti-seizure drugs**
(A) Primadone
(B) Valproic acid
(C) Ethosuccimide
(D) Embonic acid
11. **Which of the following anti-hypertensive can precipitate hypertensive crisis following abrupt cessation of therapy**
(A) Clonidine
(B) Gemfibozil
(C) Enalapril
(D) Losartan
12. **Drug of choice for the emergency control of convulsions of any type**
(A) Amiodarone
(B) Lorazepam
(C) Vigabtrin
(D) None
13. **Which of the following drug is used to prevent allograft rejection that can cause hyperlipidemia**
(A) Azathioprine
(B) Tacrolimus
(C) Sirolimus
(D) Basiliximab



MYBO's GPAT/ NIPER/ PGE CET BOOSTER PACK-8

(From MYBO Knowledge Center) : MYBO – A Voluntary Organization

14. **Reyes syndrome in children is associated with use of**
- (A) Ibuprofen
 - (B) Acetaminophen
 - (C) Celecoxib
 - (D) Aspirin
15. **A 43 year old ships pilot complains of seasonal allergies. Which of the below drug is indicated**
- (A) Cyclizine
 - (B) Doxapram
 - (C) Doxylamine
 - (D) Fexofenadine

MYBO's GPAT/ NIPER/ PGE CET BOOSTER PACK-8 is provided by

MYBO Team member:

Mr. YERNI KUMAR,

Raghu College of Pharmacy, Visakhapatnam.

NOTE:

1. **Key** for the above bits is available with MYBO's GPAT/ NIPER/ PGE CET BOOSTER PACK-9
2. If you are interested to help the GPAT/ NIPER/ PGE CET Aspirants, you can send 15 bits each with four options (correct answer underlined) to mybogroup@gmail.com After verification, we will publish on your name at www.mybogroup.com

YOUR SUCCESS IS OUR MOTTO

KEY @ MYBO's BOOSTER PACK-7

1. B	2. A	3. D	4. A	5. C
6. D	7. A	8. C	9. D	10. D
11. D	12. B	13. B	14. B	15. C